## ARE YOU READY FOR YOUR COCKROACH TREATMENT?



Proper preparation is the key to getting great results from your cockroach treatment. If you have any questions about an item listed below, don't hesitate to ask for clarification. We're here to help.

## **Preparation Steps**











## **Pre-Treatment**



Do a thorough cleaning of your kitchen and bathroom. These are locations cockroaches frequent and they will be the focus of treatment.



Remove everything from your kitchen cabinets, drawers, and under the kitchen sink. Place these items in another room, outside of the treatment area.



Remove everything from your bathroom cabinets, drawers, and under the bathroom sinks. Place these items in another room, outside of the treatment area.



Remove all garbage and recycling.



If you have pets, place them in a room that is outside of the treatment area, and close the door. This allows your representative to move freely around your home.



Cover any aquariums and turn the filtration systems off.





It is necessary that the resident/management company provide your technician with access to your home for treatment. If you're not available, please make arrangements for this access to be provided. For your safety, we request that you remain out of the home for 2 to 4 hours once access has been given.





**Post-treatment** 

Over the course of the next 7-10 days, there may be a temporary increase in activity as the treatment takes effect. Please be persistent in cleaning in order to maximize success of your service.

## **FAQ**

**Are the products you use safe?** At American Pest, we apply the methods of Integrated Pest Management, which puts an emphasis on green pest control technologies before the use of limited and targeted chemical products.

**Will my residence be cockroach-free?** Cockroaches are resilient and resourceful insects. While the cockroach treatments we perform are the most effective way to eliminate roaches, it can sometimes take more than one treatment. There is the possibility you may see a temporary increase in activity due to the disruption of their current hiding spots, this is normal. It depends on the residence and local pest pressures.